arXiv:1507.04814v1 [math.NT] 17 Jul 2015

ON DEGENERATE CARLITZ q-BERNOULLI POLYNOMIALS

TAEKYUN KIM

Abstract. In this paper, we consider the degenerate Carlitz q-Bernoulli numbers and polynomials and we investigate some properties of those polynomials.

1. Introduction

Throughout this paper, $\mathbb{Z}_p$, $\mathbb{Q}_p$ and $\mathbb{C}_p$ will, respectively, denote the ring of p-adic rational integers, the field of p-adic rational numbers and the completion of the algebraic closure of $\mathbb{Q}_p$. Let ν_p be the normalized exponential valuation of $\mathbb{C}_p$ with $p_p = p^{-\nu_p(p)} = \frac{1}{p}$. When one talks of q-extension, q is variously considered as an indeterminate, a complex number $q \in \mathbb{C}$, or p-adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, we assume that $|q| < 1$. If $q \in \mathbb{C}_p$, we assume $|q-1|_p < p^{-\frac{1}{p-1}}$ so that $q^x = \exp(x \log q)$ for $|x|_p < 1$. We use the notation $[x]_q = \frac{1-q^x}{1-q}$. Note that $\lim_{q\to 1} = x$.

In [3], L. Carlitz considered q-Bernoulli numbers as follows:

$$\beta_{0,q} = 1,\ q(q\beta_q + 1)^n - \beta_{n,q} = \begin{cases} 1, & if\ n = 1, \\ 0, & if\ n > 1, \end{cases} \tag{1.1}$$

withe the usual convention about replacing β_1^n by $\beta_{n,q}$. The q-Bernoulli polynomials are defined by

$$\beta_{n,q}(x) = \sum_{l=0}^{n} \binom{n}{l} \beta_{l,q} q^l [x]_q^{n-l},\ \text{(see [6])}. \tag{1.2}$$

In [2], L. Carlitz defined the degenerate Bernoulli polynomials which are given by the generating function to be

$$\frac{t}{(1+\lambda t)^{\frac{1}{\lambda}} - 1}(1+\lambda t)^{\frac{x}{\lambda}} = \sum_{n=0}^{\infty} \beta_n(x|\lambda)\frac{t^n}{n!},\ \text{(see [2, 3])}. \tag{1.3}$$

When $x = 0$, $\beta_n(x)(\lambda) = \beta_n(0|\lambda)$ are called the *degenerate Bernoulli numbers*. Note that $\lim_{\lambda\to 0} \beta_n(x|\lambda) = B_n(x)$, where $B_n(x)$ are ordinary Bernoulli polynomials.

Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable function on $\mathbb{Z}_p$. For $f \in UD(\mathbb{Z}_p)$, the *p-adic q-integral on $\mathbb{Z}_p$* is defined by Kim to be

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N\to\infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x,\ \text{(see [6])}. \tag{1.4}$$

The Carlitz's q-Bernoulli polynomials can be represented by p-adic q-integral on $\mathbb{Z}_p$ as follows:

$$\int_{\mathbb{Z}_p} [x+y]_q^n d\mu_q(y) = \beta_{n,q}(x),\ (n \geq 0). \tag{1.5}$$

1991 *Mathematics Subject Classification.* 05A10, 05A19.

Thus, by (1.4), we get

$$\int_{\mathbb{Z}_p} e^{[x+y]_q t} d\mu_q(y) = \sum_{n=0}^{\infty} \beta_{n,q}(x) \frac{t^n}{n!}, \text{ (see [1-8]).} \tag{1.6}$$

From (1.6), we can derive the following equation:

$$\beta_{n,q}(x) = \frac{1}{(1-q)^m} \sum_{j=0}^{m} \binom{m}{j} (-1)^j q^{ix} \frac{j+1}{[j+1]_q}, \ (m \geq 0). \tag{1.7}$$

In this paper, we consider the degenerate Carlitz q-Bernoulli polynomials and numbers and we investigate some properties of those polynomials.

2. Degenerate Carlitz q-Bernoulli numbers and polynomials

In this section, we assume that $\lambda, t \in \mathbb{C}_p$ with $|\lambda t|_p < p^{-\frac{1}{p-1}}$. In the viewpoint of (1.3), we consider the *degenerate Carlitz q-Bernoulli polynomials* which are given by the generating function to be

$$\int_{\mathbb{Z}_p} (1+\lambda t)^{\frac{1}{\lambda}[x+y]_q} d\mu_q(y) = \sum_{n=0}^{\infty} \beta_{n,q}(x|\lambda) \frac{t^n}{n!}. \tag{2.1}$$

When $x = 0$, $\beta_{n,q}(\lambda) = \beta_{n,q}(0|\lambda)$ are called the *degenerate Carlitz q-Bernoulli numbers.*

Now, we observe that

$$\begin{aligned} \int_{\mathbb{Z}_p} (1+\lambda t)^{\frac{[x+y]_q}{\lambda}} d\mu_q(y) &= \sum_{n=0}^{\infty} \int_{\mathbb{Z}_p} \binom{\frac{[x+y]_q}{\lambda}}{n} d\mu_q(y) \lambda^n t^n \\ &= \sum_{n=0}^{\infty} \int_{\mathbb{Z}_p} \left(\frac{[x+y]_q}{\lambda} \right)_n d\mu_q(y) \lambda^n \frac{t^n}{n!}, \end{aligned} \tag{2.2}$$

where $\left(\frac{[x+y]_q}{\lambda}\right)_n = \frac{[x+y]_q}{\lambda} \times \left(\frac{[x+y]_q}{\lambda} - 1\right) \times \cdots \times \left(\frac{[x+y]_q}{\lambda} - n + 1\right)$.

Now, we define $[x+y]_{n,\lambda}$ as

$$[x+y]_{n,\lambda} = [x+y]_q \left([x+y]_q - \lambda\right) \cdots \left([x+y]_q - (n-1)\lambda\right), \ (n \geq 0). \tag{2.3}$$

Therefore, by (2.1), (2.2) and (2.3), we obtain the following theorem.

Theorem 2.1. *For $n \geq 0$, we have*

$$\int_{\mathbb{Z}_p} [x+y]_{n,\lambda} d\mu_q(y) = \beta_{n,q}(x|\lambda).$$

Let $S_1(n,m)$ be the *Stirling number of the first kind* which is defined by $(x)_n = \sum_{l=0}^{n} S_1(n,l) x^l$, $(n \geq 0)$. Then, by (2.2), we get

$$\begin{aligned} \int_{\mathbb{Z}_p} \left(\frac{[x+y]_q}{\lambda} \right)_n d\mu_q(y) &= \sum_{l=0}^{n} S_1(n,l) \lambda^{-l} \int_{\mathbb{Z}_p} [x+y]_q^l d\mu_q(y) \\ &= \sum_{l=0}^{n} S_1(n,l) \lambda^{-l} \beta_{l,}(x). \end{aligned} \tag{2.4}$$

Therefore, by (2.2) and (2.4), we obtain the following theorem.

Theorem 2.2. *For $n \geq 0$, we have*

$$\beta_{n,q}(x|\lambda) = \sum_{l=0}^{n} S_1(n,l)\lambda^{n-l}\beta_{l,q}(x).$$

Note that $\lim_{\lambda\to 0}\beta_{n,q}(x|\lambda) = \beta_{n,q}(x)$.

Corollary 2.3. *For $n \geq 0$, we have*

$$\beta_{n,q}(x|\lambda) = \sum_{l=0}^{n}\sum_{j=0}^{l}\frac{S_1(n,l)}{(1-q)^l}\binom{l}{j}(-1)^j q^{jx}\frac{j+1}{[j+1]_q}\lambda^{n-l}.$$

We observe that

$$\begin{aligned}(1+\lambda t)^{\frac{[x+y]_q}{\lambda}} &= e^{\frac{[x+y]_q}{\lambda}\log(1+\lambda t)} = \sum_{n=0}^{\infty}\left(\frac{[x+y]_q}{\lambda}\right)^n\frac{1}{n!}\left(\log(1+\lambda t)\right)^n \\ &= \sum_{m=0}^{\infty}\left(\frac{[x+y]_q}{\lambda}\right)^m\frac{1}{m!}m!\sum_{n=m}^{\infty}S_1(n,m)\frac{\lambda^n t^n}{n!} \\ &= \sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}\lambda^{n-m}S_1(n,m)[x+y]_q^m\right)\frac{t^n}{n!}.\end{aligned} \tag{2.5}$$

Thus, by (2.5), we get

$$\begin{aligned}\int_{\mathbb{Z}_p}(1+\lambda t)^{\frac{[x+y]_q}{\lambda}}d\mu_q(y) &= \sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}\lambda^{n-m}S_1(n,m)\int_{\mathbb{Z}_p}[x+y]_q^m d\mu_q(x)\right)\frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}\lambda^{n-m}S_1(n,m)\beta_{m,q}(x)\right)\frac{t^n}{n!}.\end{aligned}$$

Replacing t by $\frac{1}{\lambda}\left(e^{\lambda t}-1\right)$ in (2.1), we get

$$\begin{aligned}\int_{\mathbb{Z}_p}e^{[x+y]_q t}d\mu_q(y) &= \sum_{m=0}^{\infty}\beta_{m,q}(x|\lambda)\frac{1}{m!}\frac{1}{\lambda^m}\left(e^{\lambda t}-1\right)^m \\ &= \sum_{m=0}^{\infty}\beta_{m,q}(x|\lambda)\lambda^{-m}\sum_{n=m}^{\infty}S_2(n,m)\frac{\lambda^n t^n}{n!} \\ &= \sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}\beta_{m,q}(x|\lambda)\lambda^{n-m}S_2(n,m)\right)\frac{t^n}{n!},\end{aligned} \tag{2.6}$$

where $S_2(n,m)$ is the Stirling numbers of the second kind.

We note that the left hand side of (2.6) is given by

$$\begin{aligned}\int_{\mathbb{Z}_p}e^{[x+y]_q t}d\mu_q(y) &= \sum_{n=0}^{\infty}\int_{\mathbb{Z}_p}[x+y]_q^n d\mu_q(y)\frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty}\beta_{n,q}(x)\frac{t^n}{n!}.\end{aligned} \tag{2.7}$$

Therefore, by (2.6) and (2.7), we obtain the following theorem.

Theorem 2.4. *For $n \geq 0$, we have*

$$\beta_{n,q}(x) = \sum_{m=0}^{n} \beta_{m,q}(x|\lambda)\lambda^{n-m} S_2(n,m).$$

Note that

$$\begin{aligned}(1+\lambda t)^{\frac{[x+y]_q}{\lambda}} =& (1+\lambda t)^{\frac{[x]_q}{\lambda}}(1+\lambda t)^{\frac{q^x[y]_q}{\lambda}}\\ =&\left(\sum_{m=0}^{\infty}[x]_{m,\lambda}\frac{t^m}{m!}\right)\left(\sum_{l=0}^{\infty}\frac{q^{lx}}{\lambda^l}\frac{[y]_q^l\left(\log(1+\lambda t)\right)^l}{l!}\right)\\ =&\left(\sum_{m=0}^{\infty}[x]_{m,\lambda}\frac{t^m}{m!}\right)\left(\sum_{k=0}^{\infty}\left(\sum_{l=0}^{k}\lambda^{k-l}q^{lx}[y]_q^l S_1(k,l)\right)\frac{t^k}{k!}\right)\\ =&\sum_{n=0}^{\infty}\left(\sum_{k=0}^{n}\sum_{l=0}^{k}[x]_{n-k,\lambda}\lambda^{k-l}q^{lx}[y]_q^l S_1(k,l)\binom{n}{k}\right)\frac{t^n}{n!}.\end{aligned} \tag{2.8}$$

Thus, by (2.8), we get

$$\begin{aligned}\sum_{n=0}^{\infty}\beta_{n,q}(x|\lambda)\frac{t^n}{n!} =& \sum_{n=0}^{\infty}\left(\sum_{k=0}^{n}\sum_{l=0}^{k}[x]_{n-k,\lambda}\lambda^{k-l}q^{lx}\int_{\mathbb{Z}_p}[y]_q^l d\mu_q(y) S_1(k,l)\binom{n}{k}\right)\frac{t^n}{n!}\\ =&\sum_{n=0}^{\infty}\left(\sum_{k=0}^{n}\sum_{l=0}^{k}\binom{n}{k}[x]_{n-k,\lambda}\lambda^{k-l}q^{lx}\beta_{l,q}S_1(k,l)\right)\frac{t^n}{n!}.\end{aligned} \tag{2.9}$$

Therefore, by (2.3), we obtain the following theorem.

Theorem 2.5. *For $n \geq 0$, we have*

$$\beta_{n,q}(x|\lambda) = \sum_{k=0}^{n}\sum_{l=0}^{k}\binom{n}{k}[x]_{n-k,\lambda}\lambda^{k-l}q^{lx}S_1(k,l)\beta_{l,q}.$$

For $r \in \mathbb{N}$, we define the *degenerate Carlitz q-Bernoulli polynomials of order r* as follows:

$$\begin{aligned}&\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(1+\lambda t)^{\frac{[x_1+\cdots+x_r+x]_q}{\lambda}}d\mu_1(x_1)\cdots d\mu_q(x_r)\\ =&\sum_{n=0}^{\infty}\beta_{n,q}^{(r)}(x|\lambda)\frac{t^n}{n!}.\end{aligned} \tag{2.10}$$

We observe that

$$\begin{aligned}
&\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(1+\lambda t)^{\frac{[x_1+\cdots+x_r+x]}{\lambda}}d\mu_q(x_1)\cdots d\mu_q(x_r)\\
=&\sum_{m=0}^{\infty}\lambda^{-m}\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}[x_1+\cdots+x_r+x]_q^m d\mu_q(x_1)\cdots d\mu_q(x_r)\frac{1}{m!}\left(\log(1+\lambda t)\right)^m\\
=&\sum_{m=0}^{\infty}\beta_{m,q}^{(r)}(x)\lambda^{-m}\sum_{n=m}^{\infty}S_1(n,m)\frac{\lambda^n}{n!}t^n\\
=&\sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}\lambda^{n-m}\beta_{m,q}^{(r)}S_1(n,m)\right)\frac{t^n}{n!},
\end{aligned}\tag{2.11}$$

where $\beta_{m,q}^{(r)}$ are the Carlitz q-Bernoulli numbers of order r.

Therefore, by (2.10) and (2.11), we obtain the following theorem.

Theorem 2.6. *For $n \geq 0$, we have*

$$\beta_{n,q}^{(r)}=\sum_{m=0}^{n}\lambda^{n-m}\beta_{m,q}^{(r)}S_1(n,m).$$

Replacing t by $\frac{1}{\lambda}\left(e^{\lambda t}-1\right)$ in (2.10), we have

$$\begin{aligned}
&\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}e^{[x_1+\cdots+x_r+x]_q t}d\mu_q(x_1)\cdots d\mu_q(x_r)\\
=&\sum_{m=0}^{\infty}\beta_{m,q}^{(r)}(x|\lambda)\frac{1}{m!}\lambda^{-m}\left(e^{\lambda t}-1\right)^m\\
=&\sum_{m=0}^{\infty}\beta_{m,q}^{(r)}(x|\lambda)\lambda^{-m}\sum_{n=m}^{\infty}S_2(n,m)\frac{\lambda^n t^n}{n!}\\
=&\sum_{n=0}^{\infty}\left(\sum_{m=0}^{n}\lambda^{n-m}\beta_{m,q}^{(r)}(x|\lambda)S_2(n,m)\right)\frac{t^n}{n!}.
\end{aligned}\tag{2.12}$$

The left hand side of (2.12) is given by

$$\begin{aligned}
&\int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}e^{[x_1+\cdots+x_r+x]_q t}d\mu_q(x_1)\cdots d\mu_q(x_r)\\
=&\sum_{n=0}^{\infty}\beta_{n,q}^{(r)}(x)\frac{t^n}{n!}.
\end{aligned}\tag{2.13}$$

By comparing the coefficients on the both sides of (2.12) and (2.13), we obtain the following theorem.

Theorem 2.7. *For $n \geq 0$, we have*

$$\beta_{m,q}^{(r)}(x)=\sum_{m=0}^{n}\lambda^{n-m}S_2(n,m)\beta_{m,q}^{(r)}(x|\lambda).$$

We recall that

$$\begin{aligned}\int_{\mathbb{Z}_p} f(x)d\mu_q(x) &= \lim_{N\to\infty}\frac{1}{[p^N]_q}\sum_{x=0}^{p^N-1} f(x)q^x \\ &= \lim_{N\to\infty}\frac{1}{[dp^N]_q}\sum_{x=0}^{dp^N-1} f(x)q^x,\end{aligned}$$

where $d \in \mathbb{N}$ and $f \in UD(\mathbb{Z}_p)$.

Now, we observe that

$$\beta_{n,q}(x|\lambda) = \sum_{l=0}^{n} S_1(n,l)\lambda^{n-l}\int_{\mathbb{Z}_p}[x+y]_q^l d\mu_q(y), \tag{2.14}$$

and

$$\begin{aligned}\int_{\mathbb{Z}_p}[x+y]_q^l d\mu_q(y) =&\frac{1}{[m]_q}\sum_{i=0}^{m-1} q^i[m]_q^l\int_{\mathbb{Z}_p}\left[\frac{x+i}{m}+y\right]_{q^m}^l d\mu_{q^m}(y) \\ =&[m]_q^{l-1}\sum_{i=0}^{m-1} q^i\beta_{l,q^m}\left(\frac{x+i}{m}\right),\end{aligned} \tag{2.15}$$

where $l \in \mathbb{Z}_{\geq 0}$ and $m \in \mathbb{N}$.

Therefore, by (2.14) and (2.15), we obtain the following theorem.

Theorem 2.8. *For $n \geq \mathbb{N}_{\geq 0}$, $m \in \mathbb{N}$, we have*

$$\beta_{n,q}(x|\lambda) = \sum_{l=0}^{n}\sum_{i=0}^{m-1} S_1(n,l)\lambda^{n-l}[m]_q^{l-1}q^i\beta_{l,q^m}\left(\frac{x+i}{m}\right).$$

From (1.4), we note that

$$qI_q(f_1) - I_q(f) = (q-1)f(0) + \frac{q-1}{\log q}f^{'}(0), \tag{2.16}$$

where $f^{'}(0) = \frac{df(x)}{dx}\Big|_{x=0}$.

By (2.16), we get

$$q\beta_{n,q}(x+1|\lambda) - \beta_{n,q}(x|\lambda) = (q-1)\lambda^n\left(\frac{[x]_q}{\lambda}\right)_n + \sum_{l=1}^{n-1} S_1(n,l)\lambda^{n-l}l[x]_q^{l-1}q^x, \tag{2.17}$$

where $n \in \mathbb{N}$.

Therefore, by (2.17), we obtain the following theorem.

Theorem 2.9. *For $n \geq 0$, we have*

$$q\beta_{n,q}(x+1|\lambda) - \beta_{n,q}(x|\lambda) = (q-1)\lambda^n\left(\frac{[x]_q}{\lambda}\right)_n + \sum_{l=1}^{n-1} S_1(n,l)\lambda^{n-l}l[x]_q^{l-1}q^x.$$

REFERENCES

[1] A. Bayad, J. Chikhi, *Apostol-Euler polynomials and asymptotics for negative binomial reciprocals*, Adv. Stud. Contemp. Math., **24** (2014), no. 1, 33-37.

[2] J. Choi, T. Kim, Y. H. Kim, *A note on the extended q-Bernoulli numbers and polynomials*, Adv. Stud. Contemp. Math., **21** (2011), no. 4, 351-354.

[3] L. Carlitz, *Degenerate Stirling, Bernoulli and Eulerian numbers*, Utilitas Math., **15** (1979), 51-88.

[4] L. Carlitz, *A degenerate Staudt-Clausen theorem*, Arch. Math. (Basel), **7** (1956), 28-33.

[5] D. Kang, S. J. Lee, J.-W. Park, S.-H. Rim, *On the twisted weak weight q-Bernoulli polynomials and numbers*, Proc. Jangjeon Math. Soc., **16** (2013), no. 2, 195-201.

[6] T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys., **9** (2002), no. 3, 288-299.

[7] J. W. Park, *New approach to q-Bernoulli polynomials with weight or weak weight*, Adv. Stud. Contemp. Math., **24** (2014), no. 1, 39-44.

[8] J.-J. Seo, S.-H. Rim, S.-H. Lee, D. V. Dolgy, T. Kim, *q-Bernoulli numbers and polynomials related to p-adic invariant integral on $\mathbb{Z}_p$*, Proc. Jangjeon Math. Soc., **16** (2013), no. 3, 321-326.

DEPARTMENT OF MATHEMATICS, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA.

E-mail address: `tkkim@kw.ac.kr`